

# LAMP1 Knockdown HeLa Cell Line, Heterozygous

**Catalog No.:** RM01895

## Basic Information

### Catalog No.

RM01895

### Category

Cell Line

### Parental Cell line

HeLa

### Genotype

Knockdown

## Gene Information

### Gene Symbol

LAMP1

### Species

Human

### Gene ID

3916

### Swiss Prot

P11279

### Synonyms

CD107a; LAMPA; LGP120

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

The protein encoded by this gene is a member of a family of membrane glycoproteins. This glycoprotein provides selectins with carbohydrate ligands. It may also play a role in tumor cell metastasis. [provided by RefSeq, Jul 2008]

## Product Information

### Description

LAMP1 Knockdown HeLa Cell Line is engineered from HeLa cell line with Gene-Editing Technology.

Allele-1:31bp deletion and 1bp insertion in exon3

Allele-2:62bp deletion in exon3

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

### Packaging

1 vial parental cell line and 1 vial knockout cell line

### Shipping Conditions

Dry ice

### Amount

1~5x10<sup>6</sup> cells/vial

### Storage

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

### Protocol

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO<sub>2</sub> condition.

1. Thaw the vial in 37°C water bath, and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37°C incubator with 5% CO<sub>2</sub>.
7. A subcultivation ratio of 1:2-1:4 is recommended.

## Sequencing data

---

WT    AAGCAATCACGAG\*\*\*\*\*GAGCTGCGGTTGAG  
Mut    AAGCAATCACGAG\*\*\*Deletion\*\*\*GAGCTGCGGTTGAG  
Allele-1: 31bp deletion and 1bp insertion in exon3  
  
WT    AAGCAATCACGAGA\*\*\*\*\*TGGCAGGTCAAAGG  
Mut    AAGCAATCACGAGA\*\*\*Deletion\*\*\*TGGCAGGTCAAAGG  
Allele-2: 62bp deletion in exon3

Genome sequence analysis of PCR products from parental (WT) and LAMP1 Knockdown (KD) HeLa cells, using sanger sequencing.