

AMOT Knockout 293T Cell Line, Homozygous

Catalog No.: RM01932

Basic Information

Catalog No.

RM01932

Category

Cell Line

Parental Cell line

293T

Genotype

Knockout

Gene Information

Gene Symbol

AMOT

Species

Human

Gene ID

154796

Swiss Prot

Q4VCS5

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

This gene belongs to the motin family of angiostatin binding proteins characterized by conserved coiled-coil domains and C-terminal PDZ binding motifs. The encoded protein is expressed predominantly in endothelial cells of capillaries as well as larger vessels of the placenta where it may mediate the inhibitory effect of angiostatin on tube formation and the migration of endothelial cells toward growth factors during the formation of new blood vessels. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008]

Product Information

Description

AMOT Knockout 293T Cell Line is engineered from 293T cell line with Gene-Editing Technology.

Allele-1:1bp insertion in exon1

Allele-2:1bp insertion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions

Dry ice

Amount

1~5x10⁶ cells/vial

Storage

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

Protocol

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO₂ condition.

1. Thaw the vial in 37°C water bath, and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37°C incubator with 5% CO₂.
7. A subcultivation ratio of 1:2-1:4 is recommended.

Sequencing data

WT GATGGTTGAGATCCTCTCA -GACGAGAACCGGAACTTGA
Mut GATGGTTGAGATCCTCTCA **A**GACGAGAACCGGAACTTGA
Allele-1: 1bp insertion in exon1

WT GATGGTTGAGATCCTCTCA -GACGAGAACCGGAACTTGA
Mut GATGGTTGAGATCCTCTCA **A**GACGAGAACCGGAACTTGA
Allele-2: 1bp insertion in exon1

Genome sequence analysis of PCR products from parental (WT) and AMOT knockout (KO) 293T cells, using sanger sequencing.