STAT5B Knockout HeLa Cell Line, Homozygous

Catalog No.: RM01942



Basic Information

Catalog No. RM01942

Category Cell Line

Parental Cell line HeLa

Genotype Knockout

Gene Information

Gene Symbol STAT5B

Species Human

Gene ID 6777

Swiss Prot P51692

Synonyms STAT5

Contact

6	400-999-6126
\times	cn.market@abclonal.com.cn
€	www.abclonal.com.cn

Background

The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein mediates the signal transduction triggered by various cell ligands, such as IL2, IL4, CSF1, and different growth hormones. It has been shown to be involved in diverse biological processes, such as TCR signaling, apoptosis, adult mammary gland development, and sexual dimorphism of liver gene expression. This gene was found to fuse to retinoic acid receptor-alpha (RARA) gene in a small subset of acute promyelocytic leukemias (APLL). The dysregulation of the signaling pathways mediated by this protein may be the cause of the APLL. [provided by RefSeq, Jul 2008]

Product Information

Description

STAT5B Knockout HeLa Cell Line is engineered from HeLa cell line with Gene-Editing Technology. Allele-1:exon2 was deleted

Allele-2:exon2 was deleted

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions Dry ice

Amount 1~5x10⁶ cells/vial

Storage

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

Protocol

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37° C with 5% CO₂ condition.

- 1. Thaw the vial in 37°C water bath ,and shake it to melt as soon as possible.
- 2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
- 3. Remove and discard the supernatant.
- 4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
- 5. Add 8-10mL of complete medium.
- 6. Incubate the culture at 37°C incubator with 5% CO_2 .
- 7. A subcultivation ratio of 1:2-1:4 is recommended.

Sequencing data

WT GAGGCTGTATTTAT*****GGGGGGTCTTCTTCC Mut GAGGCTGTATTTAT***Deletion***GGGGGTCTTCTTCC Allele-1: exon2 was deleted

WT GAGGCTGTATTTAT*****GGGGGGGTCTTCTTCC Mut GAGGCTGTATTTAT***Deletion***GGGGGGTCTTCTTCC

Allele-2: exon2 was deleted

Genome sequence analysis of PCR products from parental (WT) and STAT5B knockout (KO) HeLa cells, using sanger sequencing.