

PARK2 Knockout A549 Cell Lysate, Homozygous

Catalog No.: RM01970

Basic Information

Catalog No.

RM01970

Category

Cell Lysate

Parental Cell line

A549

Genotype

Knockout

Gene Information

Gene Symbol

PARK2

Species

Human

Gene ID

5071

Swiss Prot

O60260

Synonyms

AR-JP; LPRS2; PARK2; PDJ

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

The precise function of this gene is unknown; however, the encoded protein is a component of a multiprotein E3 ubiquitin ligase complex that mediates the targeting of substrate proteins for proteasomal degradation. Mutations in this gene are known to cause Parkinson disease and autosomal recessive juvenile Parkinson disease. Alternative splicing of this gene produces multiple transcript variants encoding distinct isoforms. Additional splice variants of this gene have been described but currently lack transcript support. [provided by RefSeq, Jul 2008]

Product Information

Description

PARK2 Knockout A549 Cell Line is engineered from A549 cell line with Gene-Editing technology.

Allele-1:5bp deletion in exon2

Allele-2:89bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50μL, 2μg/μL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT ATGGTTTCCAGTG*****CGATTCTGACACCA
Mut ATGGTTTCCAGTG***Deletion***CGATTCTGACACCA
Allele-1: 5bp deletion in exon2

WT ATGGTTTCCAGTG*****CGCAGGGAAGGAGC
Mut ATGGTTTCCAGTG***Deletion***CGCAGGGAAGGAGC
Allele-2: 89bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and PARK2 Knockout (KO) A549 cells, using sanger sequencing.