

# CCND1 Knockout HeLa Cell Lysate, Homozygous

Catalog No.: RM01985

## Basic Information

**Catalog No.**

RM01985

**Category**

Cell Lysate

**Parental Cell line**

HeLa

**Genotype**

Knockout

## Background

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance throughout the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK4 or CDK6, whose activity is required for cell cycle G1/S transition. This protein has been shown to interact with tumor suppressor protein Rb and the expression of this gene is regulated positively by Rb. Mutations, amplification and overexpression of this gene, which alters cell cycle progression, are observed frequently in a variety of tumors and may contribute to tumorigenesis. [provided by RefSeq, Jul 2008]

## Gene Information

**Gene Symbol**

CCND1

**Species**

Human

**Gene ID**

595

**Swiss Prot**

P24385

**Synonyms**

BCL1; D11S287E; PRAD1; U21B31

## Product Information

**Description**

CCND1 Knockout HeLa Cell Line is engineered from HeLa cell line with Gene-Editing technology.

Allele-1:29bp deletion in exon1

Allele-2:29bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles.

Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins.

Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

**Packaging**

1 vial parental cell Lysate and 1 vial knockout cell Lysate

**Shipping Conditions**

4°C

**Amount**

50μL, 2μg/μL

**Storage**

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

**Protocol**

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

## Contact

	400-999-6126
	<a href="mailto:cn.market@abclonal.com.cn">cn.market@abclonal.com.cn</a>
	<a href="http://www.abclonal.com.cn">www.abclonal.com.cn</a>

## Sequencing data

---

WT AGTGGAAACCATCC\*\*\*\*\*AACGACCGGGTGCT

Mut AGTGGAAACCATCC\*\*\*Deletion\*\*\*AACGACCGGGTGCT

Allele-1: 29bp deletion in exon1

WT AGTGGAAACCATCC\*\*\*\*\*AACGACCGGGTGCT

Mut AGTGGAAACCATCC\*\*\*Deletion\*\*\*AACGACCGGGTGCT

Allele-2: 29bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and CCND1 knockout (KO) HeLa cells, using sanger sequencing.