

# FOXM1 Knockdown HCT116 Cell Lysate, Heterozygous

**Catalog No.:** RM02034

## Basic Information

### Catalog No.

RM02034

### Category

Cell Lysate

### Parental Cell line

HCT116

### Genotype

Knockdown

## Gene Information

### Gene Symbol

FOXM1

### Species

Human

### Gene ID

2305

### Swiss Prot

Q08050

### Synonyms

FKHL16; FOXM1B; HFH-11; HFH11;  
HNF-3; INS-1; MPHOSPH2; MPP-2; MPP2;  
PIG29; TRIDENT

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

The protein encoded by this gene is a transcriptional activator involved in cell proliferation. The encoded protein is phosphorylated in M phase and regulates the expression of several cell cycle genes, such as cyclin B1 and cyclin D1. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2011]

## Product Information

### Description

FOXM1 Knockdown HCT116 Cell Line is engineered from HCT116 cell line with Gene-Editing technology.

Allele-1:105bp deletion in exon1

Allele-2:107bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles.

Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins.

Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

### Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

### Shipping Conditions

4°C

### Amount

50μL, 2μg/μL.

### Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

### Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

## Sequencing data

---

WT    CCCCTGCCCAACAG\*\*\*\*\*GTAGTGGCCATCCC  
Mut   CCCCTGCCCAACAG\*\*\*Deletion\*\*\*GTAGTGGCCATCCC  
Allele-1: 105bp deletion in exon1

WT    ATCCCCTGCCCAAC\*\*\*\*\*GTAGTGGCCATCCC  
Mut   ATCCCCTGCCCAAC\*\*\*Deletion\*\*\*GTAGTGGCCATCCC  
Allele-2: 107bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and FOXM1 Knockdown (KD) HCT116 cells, using sanger sequencing.