Leader in Biomolecular Solutions for Life Science

www.abclonal.com

ERBB2 Knockout HeLa Cell Lysate, Homozygous

Catalog No.: RM02053

Basic Information

Catalog No.
RM02053

Category
Cell Lysate

Parental Cell line
HeLa

Genotype
Knockout

Gene Information

Gene Symbol
ERBB2

Species
Human

Gene ID
2064

Swiss Prot
P04626

Synonyms
CD340; HER-2; HER-2/neu; HER2; MLN 19; NEU; NGL; TKR1

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Background

This gene encodes a member of the epidermal growth factor (EGF) receptor family of receptor tyrosine kinases. This protein has no ligand binding domain of its own and therefore cannot bind growth factors. However, it does bind tightly to other ligand-bound EGF receptor family members to form a heterodimer, stabilizing ligand binding and enhancing kinase-mediated activation of downstream signalling pathways, such as those involving mitogen-activated protein kinase and phosphatidylinositol-3 kinase. Allelic variations at amino acid positions 654 and 655 of isoform a (positions 624 and 625 of isoform b) have been reported, with the most common allele, Ile654/Ile655, shown here. Amplification and/or overexpression of this gene has been reported in numerous cancers, including breast and ovarian tumors. Alternative splicing results in several additional transcript variants, some encoding different isoforms and others that have not been fully characterized. [provided by RefSeq, Jul 2008]

Product Information

Description
ERBB2 Knockout HeLa Cell Line is engineered from HeLa cell line with Gene-Editing technology.
Allele-1:1bp insertion in exon2
Allele-2:47bp deletion in exon2
Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging
1 vial parental cell Lysate and 1 vial knockout cell Lysate


Shipping Conditions
4℃

Amount
50μL, 2μg/μL.

Storage
Store at -20℃

Protocol
To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data



Genome sequence analysis of PCR products from parental (WT) and ERBB2 knockout (KO) HeLa cells, using sanger sequencing.

For research use only. Not for therapeutic or diagnostic purposes.
Please visit http://abclonal.com for a complete listing of recommended products.