

STAT5B Knockout HeLa Cell Lysate, Homozygous

Catalog No.: RM02068

Basic Information

Catalog No.

RM02068

Category

Cell Lysate

Parental Cell line

HeLa

Genotype

Knockout

Gene Information

Gene Symbol

STAT5B

Species

Human

Gene ID

6777

Swiss Prot

P51692

Synonyms

STAT5

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Background

The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein mediates the signal transduction triggered by various cell ligands, such as IL2, IL4, CSF1, and different growth hormones. It has been shown to be involved in diverse biological processes, such as TCR signaling, apoptosis, adult mammary gland development, and sexual dimorphism of liver gene expression. This gene was found to fuse to retinoic acid receptor-alpha (RARA) gene in a small subset of acute promyelocytic leukemias (APLL). The dysregulation of the signaling pathways mediated by this protein may be the cause of the APLL. [provided by RefSeq, Jul 2008]

Product Information

Description

STAT5B Knockout HeLa Cell Line is engineered from HeLa cell line with Gene-Editing technology.

Allele-1:exon2 was deleted

Allele-2:exon2 was deleted

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50μL, 2μg/μL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT GAGGCTGTATTAT*****GGGGTCTTCTCC
Mut GAGGCTGTATTAT***Deletion***GGGGTCTTCTCC
Allele-1: exon2 was deleted

WT GAGGCTGTATTAT*****GGGGTCTTCTCC
Mut GAGGCTGTATTAT***Deletion***GGGGTCTTCTCC
Allele-2: exon2 was deleted

Genome sequence analysis of PCR products from parental (WT) and STAT5B knockout (KO) HeLa cells, using sanger sequencing.