Leader in Biomolecular Solutions for Life Science

www.abclonal.com

HMMR Knockout HeLa Cell Line, Homozygous

Catalog No.: RM02100

Basic Information

Catalog No.
RM02100

Category
Cell Line

Parental Cell line
HeLa

Genotype
Knockout

Gene Information

Gene Symbol
HMMR

Species
Human

Gene ID
3161

Swiss Prot
O75330

Synonyms
CD168; IHABP; RHAMM

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Background

The protein encoded by this gene is involved in cell motility. It is expressed in breast tissue and together with other proteins, it forms a complex with BRCA1 and BRCA2, thus is potentially associated with higher risk of breast cancer. Alternatively spliced transcript variants encoding different isoforms have been noted for this gene. [provided by RefSeq, Dec 2008]

Product Information

Description
HMMR Knockout HeLa Cell Line is engineered from HeLa cell line with Gene-Editing Technology.
Allele-1:1bp deletion in exon2
Allele-2:77bp deletion in exon2
Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging
1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions
Dry ice

Amount
1~5x10^6 cells/vial

Storage
Store at -20℃

Protocol
Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37℃ with 5% CO_2 condition.
1. Thaw the vial in 37℃ water bath ,and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37℃ incubator with 5% CO_2.
7. A subcultivation ratio of 1:2-1:4 is recommended.

Sequencing data

WT GGCTAAATGCTGCA**********TAAGGGAAAAAACA
Mut GGCTAAATGCTGCA***Deletion***TAAGGGAAAAAACA
Allele-1: 1bp deletion in exon2

WT GTTCTTCTACAGGA**********TAAGGGAAAAAACA
Mut GTTCTTCTACAGGA***Deletion***TAAGGGAAAAAACA
Allele-2: 77bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and HMMR knockout (KO) HeLa cells, using sanger sequencing.