

# **EEF2K Knockout NIH/3T3 Cell Line, Homozygous**

Catalog No.: RM02174

## **Basic Information**

### Catalog No.

RM02174

### Category

Cell Line

#### **Parental Cell line**

NIH/3T3

#### Genotype

Knockout

# **Gene Information**

### **Gene Symbol**

EEF2K

#### **Species**

Mouse

#### **Gene ID**

13631

## **Synonyms**

C86191; eEF-2K

# Contact

<u>a</u>	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

# **Background**

### **Product Information**

#### Description

EEF2K Knockout NIH/3T3 Cell Line is engineered from NIH/3T3 cell line with Gene-Editing Technology.

Allele-1:1bp deletion in exon1

Allele-2:1bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

### **Packaging**

1 vial parental cell line and 1 vial knockout cell line

# **Shipping Conditions**

**Amount** 

Dry ice

1~5x10<sup>6</sup> cells/vial

### Storage

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

#### Protoco

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at  $37^{\circ}C$  with 5% CO<sub>2</sub> condition.

- 1. Thaw the vial in 37°C water bath ,and shake it to melt as soon as possible.
- 2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
- 3. Remove and discard the supernatant.
- 4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
- 5. Add 8-10mL of complete medium.
- 6. Incubate the culture at 37°C incubator with 5% CO<sub>2</sub>.
- 7. A subcultivation ratio of 1:2-1:4 is recommended.

# Sequencing data

WT CAGAGTGCGGCTCCACAGGGTCACCAGCC
Mut CAGAGTGCGGCTCC -CAGGGTCACCAGCC

Allele-1: 1bp deletion in exon1

WT CAGAGTGCGGCTCCACAGGGTCACCAGCC
Mut CAGAGTGCGGCTCC -CAGGGTCACCAGCC

Allele-2: 1bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and EEF2K knockout (KO) NIH/3T3 cells, using sanger sequencing.