

# TDO2 Knockdown 293T Cell Lysate, Heterozygous

**Catalog No.:** RM02300

## Basic Information

### Catalog No.

RM02300

### Category

Cell Lysate

### Parental Cell line

293T

### Genotype

Knockdown

## Gene Information

### Gene Symbol

TDO2

### Species

Human

### Gene ID

6999

### Swiss Prot

P48775

### Synonyms

TDO; TO; TPH2; TRPO

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

This gene encodes a heme enzyme that plays a critical role in tryptophan metabolism by catalyzing the first and rate-limiting step of the kynurenine pathway. Increased activity of the encoded protein and subsequent kynurenine production may also play a role in cancer through the suppression of antitumor immune responses, and single nucleotide polymorphisms in this gene may be associated with autism. [provided by RefSeq, Feb 2012]

## Product Information

### Description

TDO2 Knockdown 293T Cell Line is engineered from 293T cell line with Gene-Editing technology.

Allele-1:62bp deletion in exon2

Allele-2:63bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

### Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

### Shipping Conditions

4°C

### Amount

50μL, 2μg/μL.

### Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

### Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

## Sequencing data

---

WT   AAAAACTCCCGTA\*\*\*\*\*CTATGGGAACTACC  
Mut   AAAAACTCCCGTA\*\*\*Deletion\*\*\*TATGGGAACTACC  
Allele-1: 62bp deletion in exon2  
  
WT   AAAAACTCCCGTA\*\*\*\*\*TATGGGAACTACCT  
Mut   AAAAACTCCCGTA\*\*\*Deletion\*\*\*TATGGGAACTACCT  
Allele-2: 63bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and TDO2 Knockdown (KD) 293T cells, using sanger sequencing.