Leader in Biomolecular Solutions for Life Science

www.abclonal.com

CCND3 Knockout HeLa Cell Lysate, Homozygous

Catalog No.: RM02366

Basic Information

Catalog No.
RM02366

Category
Cell Lysate

Parental Cell line
HeLa

Genotype
Knockout

Gene Information

Gene Symbol
CCND3

Species
Human

Gene ID
896

Swiss Prot
P30281

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Background

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK4 or CDK6, whose activtiy is required for cell cycle G1/S transition. This protein has been shown to interact with and be involved in the phosphorylation of tumor suppressor protein Rb. The CDK4 activity associated with this cyclin was reported to be necessary for cell cycle progression through G2 phase into mitosis after UV radiation. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2008]

Product Information

Description
CCND3 Knockout HeLa Cell Line is engineered from HeLa cell line with Gene-Editing technology.
Allele-1:86bp deletion in exon2
Allele-2:86bp deletion in exon2
Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging
1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions
4℃

Amount
50μL, 2μg/μL.

Storage
Store at -20℃

Protocol
To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

```
WT    GGAGGTGCTGGTCC**************CGACAGGCCTTGGT
Mut   GGAGGTGCTGGTCC***Deletion***CGACAGGCCTTGGT
Allele-1：86bp deletion in exon2

WT    GGAGGTGCTGGTCC**************CGACAGGCCTTGGT
Mut   GGAGGTGCTGGTCC***Deletion***CGACAGGCCTTGGT
Allele-2：86bp deletion in exon2
```

Genome sequence analysis of PCR products from parental (WT) and CCND3 knockout (KO) HeLa cells, using sanger sequencing.

For research use only. Not for therapeutic or diagnostic purposes.
Please visit http://abclonal.com for a complete listing of recommended products.