

# **GNAI3** Knockout HeLa Cell Lysate, Homozygous

Catalog No.: RM02387

# **Basic Information**

#### Catalog No.

RM02387

### Category

Cell Lysate

### **Parental Cell line**

HeLa

### Genotype

Knockout

# **Background**

Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling pathways. G proteins are composed of 3 units: alpha, beta and gamma. This gene encodes an alpha subunit and belongs to the G-alpha family. Mutation in this gene, resulting in a gly40-to-arg substitution, is associated with auriculocondylar syndrome, and shown to affect downstream targets in the G protein-coupled endothelin receptor pathway. [provided by RefSeq, Jun 2012]

# **Gene Information**

# **Gene Symbol**

**GNAI3** 

### **Species**

Human

# Gene ID

2773

### **Swiss Prot**

P08754

# Synonyms

87U6; ARCND1

### **Contact**

<u>a</u>	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

# **Product Information**

#### Description

GNAI3 Knockout HeLa Cell Line is engineered from HeLa cell line with Gene-Editing technology.

Allele-1:70bp deletion in exon1

Allele-2:71bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

## **Packaging**

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions Amount  $4^{\circ}$ C 50 $\mu$ L, 2 $\mu$ g/ $\mu$ L.

# Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

#### Protocol

To be used as WB control. Lysate is supplied in  $1\times$  SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3-5 minutes before loading onto gel.

# Sequencing data

WT TCAGCCTGCCGAGC\*\*\*\*\*\*\*\*\*\*\*\*\*GGGAGGACGGGGAA
Mut TCAGCCTGCCGAGC\*\*\*Deletion\*\*\*GGGAGGACGGGGAA
Allele-1: 70bp deletion in exon1

WT AGCCTGCCGAGCCG\*GGAGGACGGGGAAA Mut AGCCTGCCGAGCCC\*\*\*Deletion\*\*\*\*GGAGGACGGGGAAA Allele-2: 71bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and GNAI3 knockout (KO) HeLa cells, using sanger sequencing.