Leader in Biomolecular Solutions for Life Science

www.abclonal.com

# KLK3 Knockdown 293T Cell Lysate, Heterozygous

**Catalog No.: RM02401**

## Basic Information

**Catalog No.**
RM02401

**Category**
Cell Lysate

**Parental Cell line**
293T

**Genotype**
Knockdown

## Gene Information

**Gene Symbol**
KLK3

**Species**
Human

**Gene ID**
354

**Swiss Prot**
P07288

**Synonyms**
APS; KLK2A1; PSA; hK3

## Contact


400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn


## Background

Kallikreins are a subgroup of serine proteases having diverse physiological functions. Growing evidence suggests that many kallikreins are implicated in carcinogenesis and some have potential as novel cancer and other disease biomarkers. This gene is one of the fifteen kallikrein subfamily members located in a cluster on chromosome 19. Its protein product is a protease present in seminal plasma. It is thought to function normally in the liquefaction of seminal coagulum, presumably by hydrolysis of the high molecular mass seminal vesicle protein. Serum level of this protein, called PSA in the clinical setting, is useful in the diagnosis and monitoring of prostatic carcinoma. Alternate splicing of this gene generates several transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008]

## Product Information

**Description**
KLK3 Knockdown 293T Cell Line is engineered from 293T cell line with Gene-Editing technology.
Allele-1:WT
Allele-2:exon2 was deleted
Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

**Packaging**
1 vial parental cell Lysate and 1 vial knockout cell Lysate

**Shipping Conditions**
4℃

**Amount**
50μL, 2μg/μL.

**Storage**
Store at -20℃

**Protocol**
To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

## Sequencing data

WT ATCCATCTCCTATC**********CGAGTCCCCCAGTT
Mut ATCCATCTCCTATC***Deletion***CGAGTCCCCCAGTT
Allele-1: WT

WT ATCCATCTCCTATC**********GATAACCTCTAAGG
Mut ATCCATCTCCTATC***Deletion***GATAACCTCTAAGG
Allele-2: exon2 was deleted

Genome sequence analysis of PCR products from parental (WT) and KLK3 Knockdown (KD) 293T cells, using sanger sequencing.



For research use only. Not for therapeutic or diagnostic purposes.
Please visit http://abclonal.com for a complete listing of recommended products.