Leader in Biomolecular Solutions for Life Science

www.abclonal.com

# FSH Knockout 293T Cell Line, Homozygous

**Catalog No.: RM02541**

## Basic Information

**Catalog No.**
RM02541

**Category**
Cell Line

**Parental Cell line**
293T

**Genotype**
Knockout

## Gene Information

**Gene Symbol**
FSH

**Species**
Human

**Gene ID**
2488

**Swiss Prot**
P01225

**Synonyms**
HH24

## Contact


400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn


## Background

The pituitary glycoprotein hormone family includes follicle-stimulating hormone, luteinizing hormone, chorionic gonadotropin, and thyroid-stimulating hormone. All of these glycoproteins consist of an identical alpha subunit and a hormone-specific beta subunit. This gene encodes the beta subunit of follicle-stimulating hormone. In conjunction with luteinizing hormone, follicle-stimulating hormone induces egg and sperm production. Alternative splicing results in two transcript variants encoding the same protein. [provided by RefSeq, Jul 2008]

## Product Information

**Description**
FSH Knockout 293T Cell Line is engineered from 293T cell line with Gene-Editing technology.
Allele-1:158bp deletion in exon1
Allele-2:158bp deletion in exon1
Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

**Packaging**
1 vial parental cell line and 1 vial knockout cell line

**Shipping Conditions**
Dry ice

**Amount**
1~5x$10^6$ cells/vial

**Storage**
Store at -20℃

**Protocol**
Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37℃ with 5% $CO_2$ condition.

1. Thaw the vial in 37℃ water bath ,and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37℃ incubator with 5% $CO_2$.
7. A subcultivation ratio of 1:2-1:4 is recommended.

## Sequencing data

```
WT    TCCCAGACCAGGAT**********TAGGTACCATGTTT
Mut   TCCCAGACCAGGAT***Deletion***TAGGTACCATGTTT
Allele-1:   158bp deletion in exon1

WT    TCCCAGACCAGGAT**********TAGGTACCATGTTT
Mut   TCCCAGACCAGGAT***Deletion***TAGGTACCATGTTT
Allele-2:   158bp deletion in exon1
```

Genome sequence analysis of PCR products from parental (WT) and FSH knockout (KO) 293T cells, using sanger sequencing.