

# IGF1 Knockdown HeLa Cell Lysate, Heterozygous

Catalog No.: RM02550

## Basic Information

### Catalog No.

RM02550

### Category

Cell Lysate

### Parental Cell line

HeLa

### Genotype

Knockdown

## Gene Information

### Gene Symbol

IGF1

### Species

Human

### Gene ID

3479

### Swiss Prot

P05019

### Synonyms

IGF-I; IGF1; MGF

## Contact

 | 400-999-6126

 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

The protein encoded by this gene is similar to insulin in function and structure and is a member of a family of proteins involved in mediating growth and development. The encoded protein is processed from a precursor, bound by a specific receptor, and secreted. Defects in this gene are a cause of insulin-like growth factor I deficiency. Alternative splicing results in multiple transcript variants encoding different isoforms that may undergo similar processing to generate mature protein. [provided by RefSeq, Sep 2015]

## Product Information

### Description

IGF1 Knockdown HeLa Cell Line is engineered from HeLa cell line with Gene-Editing technology.

Allele-1:71bp deletion in exon2

Allele-2:WT

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

### Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

### Shipping Conditions

4°C

### Amount

50µL, 2µg/µL.

### Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

### Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

## Sequencing data

---

WT CACCATGTCCTCCT\*\*\*\*\*TGC GGGGCTGAGCT  
Mut CACCATGTCCTCCT\*\*\*Deletion\*\*\*TGC GGGGCTGAGCT  
Allele-1: 71bp deletion in exon2

WT CTGGCGCTGCGCTGCTCACCTCACCAGCTGCCACGG  
Mut CTGGCGCTGCGCTGCTCACCTCACCAGCTGCCACGG  
Allele-2: WT

Genome sequence analysis of PCR products from parental (WT) and IGF1 Knockdown (KD) HeLa cells, using sanger sequencing.