Leader in Biomolecular Solutions for Life Science

www.abclonal.com

TNFRSF18 Knockdown HeLa Cell Line, Heterozygous

Catalog No.: RM02581

Basic Information

Catalog No.
RM02581

Category
Cell Line

Parental Cell line
HeLa

Genotype
Knockdown

Gene Information

Gene Symbol
TNFRSF18

Species
Human

Gene ID
8784

Swiss Prot
Q9Y5U5

Synonyms
AITR; CD357; GITR; GITR-D

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Background

This gene encodes a member of the TNF-receptor superfamily. The encoded receptor has been shown to have increased expression upon T-cell activation, and it is thought to play a key role in dominant immunological self-tolerance maintained by CD25(+)CD4(+) regulatory T cells. Knockout studies in mice also suggest the role of this receptor is in the regulation of CD3-driven T-cell activation and programmed cell death. Three alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq, Feb 2011]

Product Information

Description
TNFRSF18 Knockdown HeLa Cell Line is engineered from HeLa cell line with Gene-Editing technology.
Allele-1:130bp deletion in exon1
Allele-2:WT
Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging
1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions
Dry ice

Amount
1~5x10^6 cells/vial

Storage
Store at -20℃

Protocol
Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37℃ with 5% CO_2 condition.
1. Thaw the vial in 37℃ water bath ,and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37℃ incubator with 5% CO_2.
7. A subcultivation ratio of 1:2-1:4 is recommended.

Sequencing data

```
WT    GCACAGCACGGGGC**************TGCCGGGTTCACAC
Mut   GCACAGCACGGGGC***Deletion***TGCCGGGTTCACAC
Allele-1:    130bp deletion in exon1

WT    GCACAGCACGGGGC**************TGCCGGGTTCACAC
Mut   GCACAGCACGGGGC**************TGCCGGGTTCACAC
Allele-2:     WT
```

Genome sequence analysis of PCR products from parental (WT) and TNFRSF18 Knockdown (KD) HeLa cells, using sanger sequencing.