ABclonal www.abclonal.com

ISL1 Knockout 293T Cell Line, Homozygous

Catalog No.: RM02674

Basic Information

Catalog No.

RM02674

Category

Cell Line

Parental Cell line

293T

Genotype

Knockout

Background

This gene encodes a member of the LIM/homeodomain family of transcription factors. The encoded protein binds to the enhancer region of the insulin gene, among others, and may play an important role in regulating insulin gene expression. The encoded protein is central to the development of pancreatic cell lineages and may also be required for motor neuron generation. Mutations in this gene have been associated with maturity-onset diabetes of the young.

Gene Information

Gene Symbol

ISL1

Species

Human

Gene ID

3670

Swiss Prot

P61371

Synonyms

Isl-1; ISLET1; Islet1

Contact

| <u>a</u> | 400-999-6126 |
|-----------|---------------------------|
| \bowtie | cn.market@abclonal.com.cn |
| • | www.abclonal.com.cn |

Product Information

Description

ISL1 Knockout cell line is engineered from 293T cell line with Gene-Editing Technology. Allele-1:exon1 was deleted

Allele-2:exon1 was deleted

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell line and 1 vial knockout cell line

Shipping Conditions

Amount

Dry ice

1~5x10⁶ cells/vial.

Storage

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

Protocol

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO_2 condition.

- 1. Thaw the vial in 37° C water bath ,and shake it to melt as soon as possible.
- 2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
- 3. Remove and discard the supernatant.
- 4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
- 5. Add 8-10mL of complete medium.
- 6. Incubate the culture at 37°C incubator with 5% CO₂.
- 7. A subcultivation ratio of 1:2-1:4 is recommended.

Sequencing data

WT GCTGTTCACCAACT*********TGCGGGGTTCTCTC
Mut GCTGTTCACCAACT***Deletion***TGCGGGGTTCTCTC
Allele-2: exon1 was deleted

Genome sequence analysis of PCR products from parental (WT) and ISL1 knockout (KO) 293T cells, using sanger sequencing.