

# NLRX1 Knockout HeLa Cell Line, Homozygous

**Catalog No.:** RM02761

## Basic Information

**Catalog No.**

RM02761

**Category**

Cell Line

**Parental Cell line**

HeLa

**Genotype**

Knockout

## Gene Information

**Gene Symbol**

NLRX1

**Species**

Human

**Gene ID**

79671

**Swiss Prot**

Q86UT6

**Synonyms**

NOD5; NOD9; NOD26; DLNB26; CLR11.3; NLRX1

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

The protein encoded by this gene is a member of the NLR family and localizes to the outer mitochondrial membrane. The encoded protein is a regulator of mitochondrial antiviral responses. Three transcript variants encoding the same protein have been found for this gene.

## Product Information

**Description**

NLRX1 Knockout cell line is engineered from HeLa cell line with Gene-Editing Technology.

Allele-1:118bp deletion in exon4

Allele-2:104bp deletion in exon4

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles.

Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins.

Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

**Packaging**

1 vial parental cell line and 1 vial knockout cell line

**Shipping Conditions**

Dry ice

**Amount**

1~5x10<sup>6</sup> cells/vial.

**Storage**

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

**Protocol**

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO<sub>2</sub> condition.

1. Thaw the vial in 37°C water bath ,and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37°C incubator with 5% CO<sub>2</sub>.
7. A subcultivation ratio of 1:2-1:4 is recommended.

## Sequencing data

---

WT CCACTCAGAAGCTA\*\*\*\*\*TACCCCTGATGAGC  
Mut CCACTCAGAAGCTA\*\*\*Deletion\*\*\*TACCCCTGATGAGC  
Allele-1: 118bp deletion in exon4

WT TATACAGCGGCACC\*\*\*\*\*AGTACCCCTGATGA  
Mut TATACAGCGGCACC\*\*\*Deletion\*\*\*AGTACCCCTGATGA  
Allele-2: 104bp deletion in exon4

Genome sequence analysis of PCR products from parental (WT) and NLRX1 knockout (KO) HeLa cells, using sanger sequencing.