

# KLRD1 Knockout 293T Cell Lysate, Homozygous

**Catalog No.:** RM50014

## Basic Information

**Catalog No.**

RM50014

**Category**

Cell Lysate

**Parental Cell line**

293T

**Genotype**

Knockout

## Gene Information

**Gene Symbol**

KLRD1

**Species**

Human

**Gene ID**

3824

**Swiss Prot**

Q13241

**Synonyms**

CD94; KLRD1

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

Natural killer (NK) cells are a distinct lineage of lymphocytes that mediate cytotoxic activity and secrete cytokines upon immune stimulation. Several genes of the C-type lectin superfamily, including members of the NKG2 family, are expressed by NK cells and may be involved in the regulation of NK cell function. KLRD1 (CD94) is an antigen preferentially expressed on NK cells and is classified as a type II membrane protein because it has an external C terminus. Several transcript variants encoding different isoforms have been found for this gene.

## Product Information

**Description**

KLRD1 Knockout cell line is engineered from 293T cell line with Gene-Editing Technology.

Allele-1:56bp deletion in exon1

Allele-2:77bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

**Packaging**

1 vial parental cell Lysate and 1 vial knockout cell Lysate

**Shipping Conditions**

4°C

**Amount**

50μL, 2μg/μL.

**Storage**

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

**Protocol**

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

## Sequencing data

---

WT TGCCAAGAAAAATG\*\*\*\*\*AAAGTCGGCATCT  
Mut TGCCAAGAAAAATG\*\*\*Deletion\*\*\*AAAGTCGGCATCT  
Allele-1: 56bp deletion in exon4

WT GACTCTGACTGCTG\*\*\*\*\*GTCGGCATCTCTGT  
Mut GACTCTGACTGCTG\*\*\*Deletion\*\*\*GTCGGCATCTCTGT  
Allele-2: 77bp deletion in exon4

Genome sequence analysis of PCR products from parental (WT) and KLRD1 knockout (KO) 293T cells, using sanger sequencing.