

RUNX1T1 Knockout 293T Cell Lysate, Homozygous

Catalog No.: RM50035

Basic Information

Catalog No.

RM50035

Category

Cell Lysate

Parental Cell line

293T

Genotype

Knockout

Gene Information

Gene Symbol

RUNX1T1

Species

Human

Gene ID

862

Swiss Prot

Q06455

SynonymsCDR; ETO; MTG8; AML1T1; ZMYND2;
CBFA2T1; AML1-MTG8; RUNX1T1

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Background

This gene encodes a member of the myeloid translocation gene family which interact with DNA-bound transcription factors and recruit a range of corepressors to facilitate transcriptional repression. The t(8;21)(q22;q22) translocation is one of the most frequent karyotypic abnormalities in acute myeloid leukemia. The translocation produces a chimeric gene made up of the 5'-region of the runt-related transcription factor 1 gene fused to the 3'-region of this gene. The chimeric protein is thought to associate with the nuclear corepressor/histone deacetylase complex to block hematopoietic differentiation. Alternative splicing results in multiple transcript variants.

Product Information

Description

RUNX1T1 Knockout cell line is engineered from 293T cell line with Gene-Editing Technology.
Allele-1:52bp deletion in exon2

Allele-2:9bp insertion and 55bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions

4°C

Amount

50μL, 2μg/μL.

Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

WT GGCATCCTCCAGAT*****AATCTAGGCTGACT
Mut GGCATCCTCCAGAT***Deletion***AATCTAGGCTGACT
Allele-1: 52bp deletion in exon2

WT CCTCCAGAT *****CTAGGCTGAC
Mut CCTCCAGATACTGTCAGC***Deletion***CTAGGCTGAC
Allele-2: 9bp Insertion and 55bp deletion in exon2

Genome sequence analysis of PCR products from parental (WT) and RUNX1T1 knockout (KO) 293T cells, using sanger sequencing.