

# DCTN4 Knockdown HeLa Cell Line, Heterozygous

Catalog No.: RM50047

## Basic Information

**Catalog No.**  
RM50047

**Category**  
Cell Line

**Parental Cell line**  
HeLa

**Genotype**  
Knockdown

## Gene Information

**Gene Symbol**  
DCTN4

**Species**  
Human

**Gene ID**  
51164

**Swiss Prot**  
Q9UJW0

**Synonyms**  
P62; DYN4

## Contact

 | 400-999-6126

 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

Enables protein N-terminus binding activity. Located in centrosome. [provided by Alliance of Genome Resources, Apr 2022]

## Product Information

### Description

DCTN4 Knockdown HeLa cell line is engineered from HeLa cell line with Gene-Editing Technology.

Allele-1:6bp deletion in exon3

Allele-2:166bp deletion in exon3

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

### Packaging

1 vial parental cell line and 1 vial knockout cell line

### Shipping Conditions

Dry ice

### Amount

1~5x10<sup>6</sup> cells/vial.

### Storage

Stored in liquid nitrogen for a long time less than -130°C. Minimizing freeze-thaw cycles.

### Protocol

Upon arrival, it should be maintained in DMEM medium with 10%(v/v) fetal bovine serum and 100U penicillin-streptomycin, at 37°C with 5% CO<sub>2</sub> condition.

1. Thaw the vial in 37°C water bath, and shake it to melt as soon as possible.
2. Transfer the cell suspension to a 15mL conical tube with pre-warmed 5mL complete medium and centrifuge 1000rpm for approximately 5 minutes at room temperature.
3. Remove and discard the supernatant.
4. Resuspend the cell pellet with 1mL pre-warmed complete medium and seed in 10cm dish.
5. Add 8-10mL of complete medium.
6. Incubate the culture at 37°C incubator with 5% CO<sub>2</sub>.
7. A subcultivation ratio of 1:2-1:4 is recommended.

## Sequencing data

---

WT CGCTGGACGTCT\*\*\*\*\*GTGGGCATGGCA  
Mut CGCTGGACGTCT\*\*Deletion\*\*GTGGGCATGGCA  
Allele-1: 6bp deletion in exon3

Genome sequence analysis of PCR products from parental (WT) and DCTN4 knockdown (KD) HeLa cells, using sanger sequencing.

WT TTGTCATGCACAG\*\*\*\*\*GACAAATCTGTAGG  
Mut TTGTCATGCACAG\*\*\*Deletion\*\*\*GACAAATCTGTAGG  
Allele-2: 166bp deletion in exon3