Leader in Biomolecular Solutions for Life Science

www.abclonal.com

MMP2 Knockout 293T Cell Lysate, Homozygous

Catalog No.: RM50086

Basic Information

Catalog No.
RM50086

Category
Cell Lysate

Parental Cell line
293T

Genotype
Knockout

Gene Information

Gene Symbol
MMP2

Species
Human

Gene ID
4313

Swiss Prot
P08253

Synonyms
CLG4; MONA; CLG4A; MMP-2; TBE-1; MMP-II; MMP2

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Background

This gene is a member of the matrix metalloproteinase (MMP) gene family, that are zinc-dependent enzymes capable of cleaving components of the extracellular matrix and molecules involved in signal transduction. The protein encoded by this gene is a gelatinase A, type IV collagenase, that contains three fibronectin type II repeats in its catalytic site that allow binding of denatured type IV and V collagen and elastin. Unlike most MMP family members, activation of this protein can occur on the cell membrane. This enzyme can be activated extracellularly by proteases, or, intracellulary by its S-glutathiolation with no requirement for proteolytical removal of the pro-domain. This protein is thought to be involved in multiple pathways including roles in the nervous system, endometrial menstrual breakdown, regulation of vascularization, and metastasis. Mutations in this gene have been associated with Winchester syndrome and Nodulosis-Arthropathy-Osteolysis (NAO) syndrome. Alternative splicing results in multiple transcript variants encoding different isoforms.

Product Information

Description
MMP2 Knockout cell line is engineered from 293T cell line with Gene-Editing Technology.
Allele-1:40bp deletion in exon1
Allele-2:40bp deletion in exon1
Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging
1 vial parental cell Lysate and 1 vial knockout cell Lysate


Shipping Conditions
4℃

Amount
50μL, 2μg/μL.

Storage
Store at -20℃

Protocol
To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data



Genome sequence analysis of PCR products from parental (WT) and MMP2 knockout (KO) 293T cells, using sanger sequencing.