

# CDX2 Knockout 293T cell lysate, Homozygous

Catalog No.: RM50180

## Basic Information

### Catalog No.

RM50180

### Category

Cell Lysate

### Parental Cell line

293T

### Genotype

Knockout

## Gene Information

### Gene Symbol

CDX2

### Species

Human

### Gene ID

1045

### Swiss Prot

Q99626

### Synonyms

CDX3; CDX-3; CDX2/AS; X2

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

This gene is a member of the caudal-related homeobox transcription factor gene family. The encoded protein is a major regulator of intestine-specific genes involved in cell growth and differentiation. This protein also plays a role in early embryonic development of the intestinal tract. Aberrant expression of this gene is associated with intestinal inflammation and tumorigenesis.

## Product Information

### Description

CDX2 Knockout cell line is engineered from 293T cell line with Gene-Editing Technology.

Allele-1:50bp deletion in exon1

Allele-2:49bp deletion in exon1

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles.

Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins.

Heterozygote: only one allele was knocked out, the mRNA transcript levels were decreased compared to wild type, and the protein expression levels were also lower than that of the wild type.

### Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

### Shipping Conditions

4°C

### Amount

50μL, 2μg/μL.

### Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

### Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

## Sequencing data

---

WT GGCCTCAACCTGGC\*\*\*\*\*ACGTGGCGGCCGCA  
Mut GGCCTCAACCTGGC\*\*\*Deletion\*\*\*ACGTGGCGGCCGCA  
Allele-1: 50bp deletion in exon1

WT GCCTCAACCTGGCG\*\*\*\*\*ACGTGGCGGCCGCA  
Mut GCCTCAACCTGGCG\*\*\*Deletion\*\*\*ACGTGGCGGCCGCA  
Allele-2: 49bp deletion in exon1

Genome sequence analysis of PCR products from parental (WT) and CDX2 knockout (KO) 293T cells, using sanger sequencing.