Leader in Biomolecular Solutions for Life Science

www.abclonal.com

MAPK1 Knockdown HeLa cell lysate, Heterozygous

Catalog No.: RM50188

Basic Information

Catalog No.
RM50188

Category
Cell Lysate

Parental Cell line
HeLa

Genotype
Knockdown

Gene Information

Gene Symbol
MAPK1

Species
Human

Gene ID
5594

Swiss Prot
P28482

Synonyms
ERK; p38; p40; p41; ERK2; ERT1; NS13; ERK-2; MAPK2; PRKM1; PRKM2; P42MAPK; p41mapk; p42-MAPK; K2

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn

Background

This gene encodes a member of the MAP kinase family. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. The activation of this kinase requires its phosphorylation by upstream kinases. Upon activation, this kinase translocates to the nucleus of the stimulated cells, where it phosphorylates nuclear targets. One study also suggests that this protein acts as a transcriptional repressor independent of its kinase activity. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions. Two alternatively spliced transcript variants encoding the same protein, but differing in the UTRs, have been reported for this gene.

Product Information

Description
MAPK1 Knockdown cell line is engineered from HeLa cell line with Gene-Editing Technology.
Allele-1:117bp deletion in exon3
Allele-2:115bp deletion in exon3
Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

Packaging
1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping Conditions
4℃

Amount
50μL, 2μg/μL.

Storage
Store at -20℃

Protocol
To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

Sequencing data

```
WT     AAACAGATATATAG**************AGCTAACGTTCTGC
Mut    AAACAGATATATAG***Deletion***AGCTAACGTTCTGC
Allele-1:   117bp deletion in exon3

WT     AAACAGATATATAG**************CTAACGTTCTGCAC
Mut    AAACAGATATATAG***Deletion***CTAACGTTCTGCAC
Allele-2:   115bp deletion in exon3
```

Genome sequence analysis of PCR products from parental (WT) and MAPK1 knockdown (KD) HeLa cells, using sanger sequencing.