# METTL3 Knockdown U2OS cell lysate, Heterozygous

Catalog No.: RM50192



## **Basic Information**

Catalog No. RM50192

Category Cell Lysate

Parental Cell line

Genotype Knockdown

## **Gene Information**

Gene Symbol METTL3

Species Human

#### Gene ID 56339

Swiss Prot Q86U44

Synonyms

M6A; IME4; Spo8; MT-A70; hMETTL3; METTL3

## Contact

6	400-999-6126
$\times$	cn.market@abclonal.com.cn
€	www.abclonal.com.cn

## Background

This gene encodes the 70 kDa subunit of MT-A which is part of N6-adenosinemethyltransferase. This enzyme is involved in the posttranscriptional methylation of internal adenosine residues in eukaryotic mRNAs, forming N6-methyladenosine.

## **Product Information**

#### Description

METTL3 Knockdown cell line is engineered from U2OS cell line with Gene-Editing Technology. Allele-1:139bp deletion in exon2

Allele-2:138bp deletion in exon2

Mammalian cells such as human, rat and mouse cells are normally diploid with two alleles. Homozygote: both alleles were knocked out, mRNA has no signal, no expression of proteins. Heterozygote: only one allele was knocked out, the mRNA transcript levels was decreased compared to wild type, and the protein expression levels was also lower than that of the wild type.

#### Packaging

1 vial parental cell Lysate and 1 vial knockout cell Lysate

Shipping	Conditions
4°C	

**Amount** 50μL, 2μg/μL.

#### Storage

Lysate is stable for 12 months when stored at -20°C. Minimizing freeze-thaw cycles.

#### Protocol

To be used as WB control. Lysate is supplied in 1× SDS sample buffer (2% SDS, 60 mM Tris-HCl pH 6.8, 10% Glycerol, 0.02% Bromophenol blue, 60 mM beta-mercaptoethanol). Lysate should be boiled for 3 - 5 minutes before loading onto gel.

## Sequencing data

WT CATTGTCTCCAACC\*\*\*\*\*\*\*\*\*\*\*CATTGCCCACTGAT Mut CATTGTCTCCAACC\*\*\*Deletion\*\*\*CATTGCCCACTGAT Allele-1: 139bp deletion in exon2

WT ATTGTCTCCAACCT\*\*\*\*\*\*\*\*\*\*CATTGCCCACTGAT Mut ATTGTCTCCAACCT\*\*\*Deletion\*\*\*CATTGCCCACTGAT Allele-2: 138bp deletion in exon2 Genome sequence analysis of PCR products from parental (WT) and METTL3 knockdown (KD) U2OS cells, using sanger sequencing.