

Rabbit anti-Human IL-7 mAb

Catalog No.: RMK0417

Basic Information

Catalog No.
RMK0417

Category
Elisa Antibody Kit

Application
ELISA

Product Information

Ig Type
Rabbit IgG

Purification
Affinity purification

Endotoxin Level

Storage
Store at -20°C.
Avoid repeated freeze-thaw cycles.

Formulation
Supplied as a 0.2um filtered solution in PBS with 0.1%Braveds MB-1,PH 7.8.

Contact

 | order@abclonal.com

 | support@abclonal.com

 | www.abclonal.com

Background

The protein encoded by this gene is a cytokine important for B and T cell development. This cytokine and the hepatocyte growth factor (HGF) form a heterodimer that functions as a pre-pro-B cell growth-stimulating factor. IL7 is found to be a cofactor for V(D)J rearrangement of the T cell receptor beta (TCRB) during early T cell development. This cytokine can be produced locally by intestinal epithelial and epithelial goblet cells, and may serve as a regulatory factor for intestinal mucosal lymphocytes. IL7 plays an essential role in lymphoid cell survival, and in the maintenance of naive and memory T cells. Alternative splicing results in multiple transcript variants encoding distinct isoforms. Additional splice variants have been described but their presence in normal tissues has not been confirmed. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection can be a potent inducer of proinflammatory cytokines and chemokines which may defend against the infection, but may also mediate destructive lung injury. Elevated seru...

Immunogen Information

Immunogen
Recombinant Human IL-7

Cross-Reactivity


/

Assay Applications


Human IL-7 Sandwich Immunoassay

		Recommended Concentration	Sample
ELISA	Capture	2-8ug/mL	Rabbit anti-Human IL-7 mAb(Cat. No. RMK0417)
	Detection	0.05-0.2ug/mL	Rabbit anti-Human IL-7 mAb (Cat. No. RMK0418)
	Standard	15.63-1000pg/mL	Recombinant IL-7 Protein


Validation Data



This standard curve is only for demonstration purposes. A standard curve should be generated for each assay.



No significant cross-reactivity or interference was observed.



Placed at 37°C for 7 days, the stability of the standard curve all conform to CV <10%.