

Catalog No.: RP00006 **Recombinant**

Species	Gene ID	Swiss Prot
Human	3606	O14116

No tag

Synonyms
IL18:|G|F:|L-18:|L-1α:|L1F4

Interleukin-18 (IL-18, also known as interferon-gamma inducing factor) is a proinflammatory cytokine that belongs to the IL-1 superfamily and is produced by macrophages and other cells. IL-18 activation is induced by infection or tissue damage and contributes to disease pathology in chronic inflammation. In the presence of IL-12 or IL-15, IL-18 enhances anti-viral Th1 immune responses by inducing IFN-gamma production and the cytolytic activity of CD8+ T cells and NK cells. In the absence of IL-12 or IL-15, however, IL-18 promotes production of the Th2 cytokines IL-4 and IL-13 by CD4+ T cells and basophils. In the presence of IL-1 beta or IL-23, IL-18 induces the antigen-independent production of IL-17 by gamma delta T cells and CD4+ T cells. IL-18 also promotes myeloid dendritic cell maturation and triggers neutrophil respiratory burst. In cancer, IL-18 exhibits diverse activities including enhancing anti-tumor immunity, inhibiting or promoting angiogenesis, and promoting tumor cell metastasis.

Source <i>E. coli</i>	Purification ≥ 95 % as determined by SDS- PAGE
---------------------------------	--

18.22 kDa 18 kDa

< 0.1 EU/μg of the protein by LAL method.

Lyophilized from a 0.22 μ m filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

Recombinant Human IL-18 Protein is produced by *E. coli* expression system. The target protein is expressed with sequence (Tyr37-Asp193) of human IL-18 (Accession #NP_001553.1).

1. Measured by its binding ability in a functional ELISA. Immobilized recombinant Human IL18 at 5 µg/mL (100µL/well) can bind recombinant Human IL18BPα with a linear range of 0.156-1.372 µg/mL. 2. Measured by its ability to induce IFN-γ secretion by KG1 human acute myelogenous leukemia cells in the presence of TNF-α (Catalog: RP00001). The ED₅₀ for this effect is 1.44-5.76 ng/mL, corresponding to a specific activity of $1.7 \times 10^5 \sim 6.9 \times 10^5$ units/mg.


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.
After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.
Avoid repeated freeze/thaw cycles.

 | 400-999-6126


 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn


Validation Data



Recombinant Human IL-18 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized recombinant Human IL18 at 5μg/mL (100μL/well) can bind recombinant Human IL18BPα with a linear range of 0.156-1.372μg/mL.



Recombinant Human IL-18 induce IFN-γ secretion by KG1 human acute myelogenous leukemia cells in the presence of TNF-α(Catalog: RP00001). The ED₅₀ for this effect is 1.44-5.76 ng/mL, corresponding to a specific activity of 1.7×10⁵~6.9×10⁵ units/mg.