

# Recombinant Human Oncostatin-M/OSM Protein

Catalog No.: RP00054 **Recombinant** **1 Publications**

## Sequence Information

Species	Gene ID	Swiss Prot
Human	5008	P13725

### Tags

No tag

### Synonyms

OSM; oncostatin-M

## Product Information

Source	Purification
<i>E. coli</i>	≥ 95% as determined by reducing SDS-PAGE.

Calculated MW	Observed MW
22.15 kDa	20-25 kDa

### Endotoxin

< 0.01 EU/μg of the protein by LAL method

### Formulation

Lyophilized from a 0.22 μm filtered solution of 20mM Tris, 150mM NaCl, 5% glycerol, pH 7.4. Contact us for customized product form or formulation.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

## Background

This protein is a member of the leukemia inhibitory factor/oncostatin-M (LIF/OSM) family. The preproprotein is proteolytically processed to generate the mature protein. This protein is a secreted cytokine and growth regulator that inhibits the proliferation of a number of tumor cell lines. This protein also regulates the production of other cytokines, including interleukin 6, granulocyte-colony stimulating factor and granulocyte-macrophage colony stimulating factor in endothelial cells.

## Basic Information

### Description

Recombinant Human Oncostatin-M/OSM Protein is produced by *E. coli* expression system. The target protein is expressed with sequence (Ala26-Arg221) of human Oncostatin-M/OSM (Accession #NP\_065391.1).

### Bio-Activity

1. At concentrations ranging from 0.01 to 5 ng/mL, M5 factor (RP03505 IL1A, RP00212 IL17A, RP01616 IL22, RP00993 TNFα, RP00054 OSM) promotes the proliferation of HaCaT cells. 2. Measured in a cell proliferation assay using TF-1 Human erythroleukemic cells. The ED<sub>50</sub> for this effect is 0.76-3.04 ng/mL, corresponding to a specific activity of 3.9×10<sup>5</sup>~1.31×10<sup>6</sup> units/mg.

### Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

### Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.


Avoid repeated freeze/thaw cycles.

### Operational Notes


For your safety and health, please wear a lab coat and disposable gloves for handling.




## Validation Data



Recombinant Human Oncostatin-M/OSM Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



At concentrations ranging from 0.01 to 5 ng/mL, M5 factor (RP03505 IL1A,RP00212 IL17A,RP01616 IL22,RP00993 TNF $\alpha$ ,RP00054 OSM)promotes the proliferation of HaCaT cells.



Measured in a cell proliferation assay using TF-1 Human erythroleukemic cells. The ED<sub>50</sub> for this effect is 0.76-3.04 ng/mL, corresponding to a specific activity of  $3.9 \times 10^5 \sim 1.31 \times 10^6$  units/mg.