Recombinant Human B7-1/CD80 Protein

ABclonal

www.abclonal.com

Catalog No.: RP00065 Recombinant

Sequence Information

Species Gene ID **Swiss Prot** Human 941 P33681

Tags

C-His

Synonyms

B7;B7-1;B7.1;BB1;CD28LG;CD28LG1;LAB 7:CD80

Product Information

Purification HEK293 cells ≥ 90 % as

determined by SDS-

PAGE.

Calculated MW Observed MW

24.71 kDa 40-50 kDa

Endotoxin

< 0.1 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

<u>a</u>	400-999-6126
\bowtie	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

Background

The protein is a membrane receptor that is activated by the binding of CD28 or CTLA-4. The activated protein induces T-cell proliferation and cytokine production. This protein can act as a receptor for adenovirus subgroup B and may play a role in lupus neuropathy.

Basic Information

Description

Recombinant Human B7-1/CD80 Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Val35-Asn242) of human B7-1/CD80 (Accession #NP 005182.1) fused with a 6×His tag at the C-terminus.

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized Recombinant human CD80 at 2 µg/mL (100 µL/well) can bind Recombinant human CTLA-4 with a linear range of 10-50 ng/mL.


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human B7-1/CD80 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized Recombinant human CD80 at 2 $\mu g/mL$ (100 $\mu L/well)$ can bind Recombinant human CTLA-4 with a linear range of 10-50 ng/mL.