

# Recombinant Human Fc-gamma RIIb/CD32b Protein

Catalog No.: RP00075 Recombinant

# **Sequence Information**

**Species** Gene ID **Swiss Prot** Human 2213 P31994

## **Tags**

C-His

### **Synonyms**

FCGR2B;CD32;CD32B;FCG2;FCGR2;IGFR

# **Product Information**

**Purification** HEK293 cells

> 97% by SDS-PAGE.

## **Endotoxin**

< 0.1 EU/µg of the protein by LAL method.

#### **Formulation**

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

## Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## **Contact**

6	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
$\odot$	www.abclonal.com.cn

# **Background**

The protein is a low affinity receptor for the Fc region of immunoglobulin gamma complexes. The protein is involved in the phagocytosis of immune complexes and in the regulation of antibody production by B-cells.

## **Basic Information**

#### Description

Recombinant Human Fc-gamma RIIb/CD32b Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Ala46-Pro217) of human Fc gamma RIIB/C (Accession #NP 003992.3) fused with a 6×His tag at the C-terminus.

### **Bio-Activity**

1. Measured by its binding ability in a functional ELISA. Immobilized Recombinant Human FCGR2B at 2 μg/mL (100 μL/well) can bind biotinylated Human IgG1 with a linear range of 0.0048-2.81 µg/mL.|2.Measured by its binding ability in a functional ELISA. Immobilized Human FCGR2B (Catalog:RP00075) at 2μg/mL (100 μL/well) can bind FCGR2B Rabbit pAb (Catalog: A12553) with a linear range of 0.3-77.48ng/mL.


#### Storage

Store the lyophilized protein at -20°C to -80 °C for long term.


After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.

Avoid repeated freeze/thaw cycles.


# **Validation Data**




Recombinant Human Fc-gamma RIIb/CD32b Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 25-35 kDa.



The purity of Human Fc gamma RIIB/C Protein (Cat.RP00075) was greater than 95% as determined by SEC-HPLC.



Immobilized Recombinant Human FCGR2B at 2 $\mu$ g/mL (100 $\mu$ L/well) can bind biotinylated Human IgG1 with a linear range of 0.0048-2.81 $\mu$ g/mL.



Immobilized Human FCGR2B (Catalog:RP00075) at 2 $\mu$ g/mL (100  $\mu$ L/well) can bind FCGR2B Rabbit pAb (Catalog: A12553) with a linear range of 0.3-77.48ng/mL.