

**Catalog No.: RP00080** **Recombinant**

Species	Gene ID	Swiss Prot
Human	201633	O495A1

## C-His

## VSIG9; VSTM3; WUCAM; TIGIT

<b>Source</b>	<b>Purification</b>
HEK293 cells	≥ 95 % as determined by SDS-PAGE

13.90 kDa                      15-20 kDa

< 0.1 EU/μg of the protein by LAL method.

Lyophilized from a 0.22  $\mu\text{m}$  filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

 | 400-999-6126

 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

This protein is a member of the PVR (poliovirus receptor) family of immunoglobulin proteins. The protein is expressed on several classes of T cells including follicular B helper T cells (TFH). The protein has been shown to bind PVR with high affinity; this binding is thought to assist interactions between TFH and dendritic cells to regulate T cell dependent B cell responses.

Recombinant Human TIGIT Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Met22-Pro141) of human TIGIT (Accession #NP\_776160.2) fused with a 6×His tag at the C-terminus.


Measured by its binding ability in a functional ELISA. Immobilized Human TIGIT at 1.5  $\mu\text{g/mL}$  (100  $\mu\text{L}$ /well) can bind Human CD155 with a linear range of 0.15-1.2  $\mu\text{g/mL}$ .

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

# Validation Data



Recombinant Human TIGIT Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized Human TIGIT at 1.5 μg/mL (100 μL/well) can bind Human CD155 with a linear range of 0.15-1.2μg/mL.