Recombinant Human FCAR/CD89 Protein

Catalog No.: RP00113 Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
Human	2204	P24071

dii 22

Tags C-hFc&His

Synonyms FCAR;CD89;CTB-61M7.2;FcalphaRI

Product Information

Source	Purification
HEK293 cells	≥ 95 % as
	determined by SDS PAGE.

Calculated MWObserved MW50.27 kDa65-75 kDa

Endotoxin

< 0.1 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

6	400-999-6126
\bowtie	cn.market@abclonal.com.cn
€	www.abclonal.com.cn

laA-or

This protein is a member of the immunoglobulin gene superfamily and receptor for the Fc region of IgA. The receptor is a transmembrane glycoprotein present on the surface of myeloid lineage cells such as neutrophils, monocytes, macrophages, and eosinophils, where it mediates immunologic responses to pathogens. It interacts with IgA-opsonized targets and triggers several immunologic defense processes, including phagocytosis, antibody-dependent cell-mediated cytotoxicity, and stimulation of the release of inflammatory mediators. Multiple alternatively spliced transcript variants encoding different isoforms have been described for this gene.

Basic Information

Description

Background

Recombinant Human FCAR/CD89 Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Gln22-Asn227) of human CD89/FCAR (Accession #NP_001991.1) fused with an Fc, 6×His tag at the C-terminus.

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized Human IGHA1 (Catalog: RPT0023) at 5 μ g/mL (100 μ L/well) can bind Human CD89/FCAR (Catalog: RP00113) with a linear range of 0.01-7.4 ng/mL.


Storage

Store at -20°C.Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.





Recombinant Human FCAR/CD89 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.

Immobilized Human IGHA1 (Catalog: RPT0023) at 5 μ g/mL (100 μ L/well) can bind Human CD89/FCAR (Catalog: RP00113) with a linear range of 0.01-7.4 ng/mL.