

Catalog No.: RP00173 **Recombinant**

Species	Gene ID	Swiss Prot
Human	3592	P29459

Tags
C-hFc&His

Synonyms
P35; CLMF; NFSK; NKSF1; IL-12A; IL12A

Source HEK293 cells	Purification ≥ 95 % as determined by SDS- PAGE
-------------------------------	--

Calculated MW	Observed MW
49.33 kDa	70 kDa

< 0.1 EU/μg of the protein by LAL method.

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Interleukin-12 subunit alpha (IL12A/IL-12p35) is also known as Cytotoxic lymphocyte maturation factor 35 kDa subunit, cytotoxic lymphocyte maturation factor 1, p35, NK cell stimulatory factor chain 1, and interleukin-12 alpha chain. IL12A/IL-12p35 is a subunit of a cytokine that acts on T and natural killer cells, and has a broad array of biological activities. The cytokine is a disulfide-linked heterodimer composed of the 35-kD subunit encoded by this gene, and a 40-kD subunit that is a member of the cytokine receptor family. This cytokine is required for the T-cell-independent induction of interferon (IFN)-gamma, and is important for the differentiation of both Th1 and Th2 cells. The responses of lymphocytes to this cytokine are mediated by the activator of transcription protein STAT4. Nitric oxide synthase 2A (NOS2A/NOS2) is found to be required for the signaling process of this cytokine in innate immunity.


Recombinant Human IL-12A/IL-12 p35 Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Arg23-Ser219) of human IL12A/NKSF1 (Accession #NP_002178.2) fused with an Fc, 6xHis tag at the C-terminus.

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human IL-12A/IL-12 p35
Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.