

Recombinant Human IL-6 Protein

Catalog No.: RP00201 Recombinant 4 Publications

Sequence Information

Species **Gene ID** **Swiss Prot**
 Human 3569 P05231

Tags

C-His

Synonyms

IL6;BSF-2;BSF2;CDF;HGF;HSF;IFN-
 beta-2;IFNB2;IL-6

Background

Interleukin-6 (IL-6) is a multifunctional α -helical cytokine that regulates cell growth and differentiation of various tissues, which is known particularly for its role in the immune response and acute phase reactions. The encoded protein has been shown to be an endogenous pyrogen capable of inducing fever in people with autoimmune diseases or infections. The protein is primarily produced at sites of acute and chronic inflammation, where it is secreted into the serum and induces a transcriptional inflammatory response through interleukin 6 receptor, alpha. The functioning of this protein is implicated in a wide variety of inflammation-associated disease states, including susceptibility to diabetes mellitus and systemic juvenile rheumatoid arthritis.

Basic Information

Description

Recombinant Human IL-6 Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Val30-Met212) of human IL6 (Accession #NP_000591.1) fused with a 6xHis tag at the C-terminus.

Bio-Activity

1. Measured by its binding ability in a functional ELISA. Immobilized Human IL6R at 1 μ g/mL (100 μ L/well) can bind Human IL-6 with a linear range of 2-15 ng/mL. 2. Measured in a cell proliferation assay using TF-1 Human erythroleukemic cells. The ED_{50} for this effect is 0.10-0.40 ng/mL. The specific activity of Recombinant Human IL-6 is $>1.2 \times 10^8$ IU/mg, which is calibrated against the human IL-6 WHO International Standard (NIBSC code: 21/308).

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.

Product Information

Source **Purification**
 HEK293 cells $\geq 95\%$ as
 determined by SDS-PAGE.

Calculated MW **Observed MW**
 21.65 kDa 23-30

Endotoxin
 < 0.1 EU/ μ g of the protein by LAL method.

Formulation
 Lyophilized from a 0.22 μ m filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.


Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.


Contact

	400-999-6126
	cn.market@abclonal.com.cn
	www.abclonal.com.cn


Validation Data



Recombinant Human IL-6 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized Human IL6R at 1 μ g/mL (100 μ L/well) can bind Human IL-6 with a linear range of 2-15 ng/mL.



Measured in a cell proliferation assay using TF-1 Human erythroleukemic cells. The ED_{50} for this effect is 0.10-0.40 ng/mL. The specific activity of Recombinant Human IL-6 is $>1.2 \times 10^8$ IU/mg, which is calibrated against the human IL-6 WHO International Standard (NIBSC code: 21/308).