

www.abclonal.com

ABclonal

Recombinant Human TACTILE/CD96 Protein

Catalog No.: RP00346 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 10225 P40200-2

Tags

C-mFc

Synonyms

CD96 molecule; CD96; DKFZp667E2122; TACTILE

Product Information

Source Purification HEK293 cells ≥ 95 % as

determined by SDS-

PAGE.

Calculated MW Observed MW

79.8 kDa 120-160 kDa

Endotoxin

< 1 EU/µg of the protein by LAL method.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

<u>a</u>	400-999-6126
\bowtie	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

Background

The receptors CD96 and TIGIT are expressed on the surface of T and natural killer (NK) cells, and recent studies suggest both play important inhibitory roles in immune function. CD96 has been shown to modulate immune cell activity in mice, with Cd96-/-mice displaying hypersensitive NK-cell responses to immune challenge and significant tumor resistance. The counterbalance between the putative inhibitory CD96 and TIGIT receptors and the activating receptor, CD226, offers unique strategies for immuno-oncology drug development.

Basic Information

Description

Recombinant Human TACTILE/CD96 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Val22-Met503) of Human TACTILE/CD96 (Accession #P40200-2) fused with a C-mFc tag at the C-terminus.

Bio-Activity

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.


Avoid repeated freeze/thaw cycles.

Operational Notes


For your safety and health, please wear a lab coat and disposable gloves for handling.

^{*} DODOODOODOODOODOO


Validation Data



Recombinant Human TACTILE/CD96 Protein was determined by Tris-Bis PAGE under reducing conditions.



Immobilized Human CD96, mFc Tag at 5 $\mu g/mL$ (100 $\mu L/Well)$ on the plate. Dose response curve for Human CD155, hFc Tag with the EC $_{50}$ of 0.21 $\mu g/mL$ determined by ELISA.



Immobilized Human CD96, mFc Tag at 2 μ g/mL (100 μ L/Well) on the plate. Dose response curve for Anti-CD96 Antibody, hFc Tag with the EC $_{50}$ of 31.8ng/mL determined by ELISA (QC Test).