

Catalog No.: RP00346 **Recombinant**

Species	Gene ID	Swiss Prot
Human	10225	P40200-2

C-mFc

CD96 molecule; CD96; DKFZp667E2122;
TACTILE

Source	Purification
HEK293 cells	≥ 95 % as determined by SDS-PAGE.

79.8 kDa 120-160 kDa

< 1 EU/μg of the protein by LAL method.

Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

The receptors CD96 and TIGIT are expressed on the surface of T and natural killer (NK) cells, and recent studies suggest both play important inhibitory roles in immune function. CD96 has been shown to modulate immune cell activity in mice, with Cd96-/- mice displaying hypersensitive NK-cell responses to immune challenge and significant tumor resistance. The counterbalance between the putative inhibitory CD96 and TIGIT receptors and the activating receptor, CD226, offers unique strategies for immunoncology drug development.


Recombinant Human TACTILE/CD96 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Val22-Met503) of Human TACTILE/CD96 (Accession #P40200-2) fused with a C-mFc tag at the C-terminus.

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Validation Data



Recombinant Human TACTILE/CD96 Protein was determined by Tris-Bis PAGE under reducing conditions.



Immobilized Human CD96, mFc Tag at 5 µg/mL (100 µL/Well) on the plate. Dose response curve for Human CD155, hFc Tag with the EC₅₀ of 0.21 µg/mL determined by ELISA.



Immobilized Human CD96, mFc Tag at 2 µg/mL (100 µL/Well) on the plate. Dose response curve for Anti-CD96 Antibody, hFc Tag with the EC₅₀ of 31.8ng/mL determined by ELISA (QC Test).