

Biotinylated Recombinant Human IL-15RA/CD215 Protein

Catalog No.: RP00436B **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	3601	Q13261-1

Tags

C-His&Avi

Synonyms

IL-15 R alpha; CD215; IL-15RA;
MGC104179

Product Information

Source

HEK293 cells

Purification

≥ 95 % as
determined by SDS-
PAGE; ≥ 95 % as
determined by
HPLC.

Calculated MW

21.3 kDa

Observed MW

50-70 kDa

Endotoxin

< 1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from 0.22 μm filtered
solution in PBS (pH 7.4). Normally 8%
trehalose is added as protectant before
lyophilization.

Reconstitution

Centrifuge the vial before opening.
Reconstitute to a concentration of
0.1-0.5 mg/mL in sterile distilled water.
Avoid vortex or vigorously pipetting the
protein. For long term storage, it is
recommended to add a carrier protein or
stabilizer (e.g. 0.1% BSA, 5% HSA, 10%
FBS or 5% Trehalose), and aliquot the
reconstituted protein solution to
minimize free-thaw cycles.

Background

Interleukin-15 receptor alpha (IL-15R alpha) is a high affinity IL-15 binding protein that is crucial for mediating IL-15 functions such as memory CD8 T cell proliferation and NK, NK/T cell, and intestinal intraepithelial lymphocyte development.

Basic Information

Description

Biotinylated Recombinant Human IL-15RA/CD215 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ile31-Thr205) of Human IL-15RA/IL-15 R alpha/CD215 (Accession #Q13261-1) fused with a C-His&Avi tag at the C-terminus.

Bio-Activity

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.
After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.
Avoid repeated freeze/thaw cycles.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn




Validation Data



Biotinylated Recombinant Human IL-15RA/CD215 Protein was determined by Tris-Bis PAGE under reducing conditions.



The purity of Biotinylated Human IL-15RA is greater than 95% as determined by SEC-HPLC.



Immobilized Human IL-15 at 2 µg/mL (100 µL/Well) on the plate. Dose response curve for Biotinylated Human IL-15RA, His Tag with the EC₅₀ of 55.8ng/mL determined by ELISA.