

Biotinylated Recombinant Human Fc-gamma RIII beta/FCGR3B/CD16b Protein

Catalog No.: RP00471B Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 2215 075015-1

Tags C-His&Avi

Synonyms

CD16; Fc gamma RIIIB; FCG3; FCGR3B; FcgRIIIB; FCRIIIB; IGFR3; CD16b (NA2); CD16B; FCG3B; FCG3; FCGR3; FCR-10

Product Information

Source Purification HEK293 cells ≥ 95 % as

determined by SDS-PAGE;≥ 95 % as determined by HPLC.

Calculated MW Observed MW

23.7 kDa 47-53 kDa

Endotoxin

< 1 EU/ μg of the protein by LAL method.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

2	400-999-6126
\times	cn.market@abclonal.com.cn

Background

Human Fc gamma RIIIB/CD16b Protein is a receptor for the Fc region of immunoglobulins gamma. Low affinity receptor. Binds complexed or aggregated IgG and also monomeric IgG. Contrary to III-A, is not capable to mediate antibody-dependent cytotoxicity and phagocytosis. May serve as a trap for immune complexes in the peripheral circulation which does not activate neutrophils.

Basic Information

Description

Biotinylated Recombinant Human Fc-gamma RIII beta/FCGR3B/CD16b Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gly17-Ser200) of Human Fc-gamma RIII beta/FCGR3B/CD16b (Accession #075015-1) fused with a C-His&Avi tag at the C-terminus.

Bio-Activity

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Validation Data



Biotinylated Recombinant Human Fc-gamma RIII beta/FCGR3B/CD16b Protein was determined by Tris-Bis PAGE under reducing conditions.



The purity of Biotinylated Human Fc gamma RIIIB (NA2) is greater than 95% as determined by SEC-HPLC.



Biotinylated Human Fc gamma RIIIB (NA2), His Tag captured on CM5 Chip via Anti-his antibody can bind Rituximab with an affinity constant of 3.50 μM as determined in SPR assay (Biacore T200).