

Recombinant Mouse Fc-gamma RIII/CD16 Protein

Catalog No.: RP00761 **Recombinant** **1 Publications**

Sequence Information

Species	Gene ID	Swiss Prot
Mouse		P08508

Tags

C-His

Synonyms

Low affinity immunoglobulin gamma Fc region receptor III; Fcgr3; Fc gamma Receptor III; CD-antigen 16; CD16; FcRIII; IgG Fc receptor III

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

Endotoxin

< 1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.2 μm filtered solution of PBS, pH7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

Background

Low affinity immunoglobulin gamma Fc region receptor III (Fc gamma RIII/CD16) is a member of the Ig superfamily. Based on close relationships in their extracellular domains, the Fc gamma Rs have been divided into three classes composing of Fc gamma RI (CD64), Fc gamma RII (CD32), and Fc gamma RIII (CD16). Each group may be encoded by multiple genes and exist in different isoforms depending on species and cell type. Mouse CD16 is a type I transmembrane protein having two extracellular Ig-like domains consisting of immunoglobulin domain, repeat, signal and transmembrane, transmembrane helix. It is expressed on a variety of myeloid and lymphoid cells and associates with Fc R gamma to deliver an activating signal upon ligand binding. Fcgr3 is IgG binding and activation or inhibition of immune responses such as antibody-dependent cellular cytotoxicity, phagocytosis, cell surface receptor signaling pathway and positive regulation of type I/II/III hypersensitivity.

Basic Information

Description

Recombinant Mouse Fc-gamma RIII/CD16 Protein is produced by Human cells expression system. The target protein is expressed with sequence (Ala31-Thr215) of mouse Fc γ RIIIA/FCGR3/CD16 (Accession #P08508) fused with a 6×His tag at the C-terminus.

Bio-Activity

Storage

Store the lyophilized protein at -20 °C to -80 °C for long term. After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week. Avoid repeated freeze/thaw cycles.


Contact

☎		400-999-6126
---	--	--------------

✉		cn.market@abclonal.com.cn
---	--	---------------------------

🌐		www.abclonal.com.cn
---	--	---------------------

Validation Data



Recombinant Mouse Fc-gamma RIII/CD16
Protein was determined by SDS-PAGE under
reducing conditions with Coomassie Blue.