

# Recombinant Human CD44 Protein

Catalog No.: RP00973

Recombinant

2 Publications

## Sequence Information

Species	Gene ID	Swiss Prot
Human	960	P16070-1

### Tags

C-hFc&amp;His

### Synonyms

CD44; CDW44; CSPG8; ECMR-III; HCELL; HUTCH-I; IN; LHR; MC56; MDU2; MDU3; MIC4; Pgp1; CD44 antigen; CDW44; CSPG8; ECMR-III; HCELL; HUTCH-I; IN; LHR; MC56; MDU2; MDU3; MIC4; Pgp1; CD44

## Product Information

### Source

HEK293 cells

### Purification

≥ 95 % as determined by SDS-PAGE.

### Calculated MW Observed MW

48.86 kDa

70-85 kDa

### Endotoxin

&lt; 1 EU/μg of the protein by LAL method.

### Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

## Background

This protein is a cell-surface glycoprotein involved in cell-cell interactions, cell adhesion and migration. It is a receptor for hyaluronic acid (HA) and can also interact with other ligands, such as osteopontin, collagens, and matrix metalloproteinases (MMPs). This protein participates in a wide variety of cellular functions including lymphocyte activation, recirculation and homing, hematopoiesis, and tumor metastasis. Transcripts for this gene undergo complex alternative splicing that results in many functionally distinct isoforms, however, the full length nature of some of these variants has not been determined. Alternative splicing is the basis for the structural and functional diversity of this protein, and may be related to tumor metastasis.

## Basic Information

### Description

Recombinant Human CD44 Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Gln21-Pro220) of human CD44 (Accession #NP\_000601.3) fused with an Fc, 6xHis tag at the C-terminus.

### Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized Hyaluronic Acid at 100 μg/mL (100 μL/well) can bind Human CD44 with a linear range of 20-330 ng/mL.

### Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


## Contact

☎ | 400-999-6126


✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)



Validation Data



Recombinant Human CD44 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized Hyaluronic Acid at 100 µg/mL (100 µL/well) can bind Recombinant Human CD44 with a linear range of 20-330 ng/mL.