

Recombinant Human BTLA/CD272 Protein

Catalog No.: RP01052 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	151888	Q7Z6A9-2

Tags

C-hFc&His

Synonyms

BTLA;BTLA1;CD272

Product Information

Source	Purification
HEK293 cells	> 90% by SDS-PAGE.

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

Background

Basic Information

Description

Active Recombinant Human BTLA Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Lys31-Thr134) of human BTLA (Accession #NP_001078826.1.) fused with an Fc, 6xHis tag at the C-terminus.

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized Recombinant human BTLA at 3 μg/mL (100 μL/well) can bind Biotinylated Recombinant human HVEM with a linear range of 18-72 ng/mL.

Storage

Store the lyophilized protein at -20°C to -80 °C for long term.

After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Contact

 | 400-999-6126


 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Validation Data



Active Active Recombinant Human BTLA Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 50-60kDa.



Immobilized Recombinant human BTLA at 3µg/mL (100 µL/well) can bind Biotinylated Recombinant human HVEM with a linear range of 18-72 ng/mL.