

Recombinant Human MERTK Protein

Catalog No.: RP01198 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 10461 Q12866

Tags

C-His

Synonyms

MER; RP38; Tyro12; c-Eyk; cmer;MERTK;RP38;Tyro12;c-Eyk;c-mer

Product Information

Source Purification HEK293 cells ≥ 95 % as

determined by SDS-

PAGE.

Calculated MW Observed MW

52.34 kDa 70-120 kDa

Endotoxin

< 0.1 EU/ μ g of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

<u>a</u>		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

Background

Basic Information

Description

Recombinant Human MERTK Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Arg26-Ala499) of human Mer/MERTK (Accession #NP_006334.2) fused with a 6×His tag at the C-terminus.

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized Human MERTK at 1 μ g/mL (100 μ L/well) can bind MERTK Rabbit mAb with a linear range of 0.2-1.2 ng/mL.

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.

Validation Data



Recombinant Human MERTK Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized Human MERTK at $1\mu g/mL$ (100 $\mu L/well)$ can bind MERTK Rabbit mAb with a linear range of 0.2-1.2ng/mL.