

Recombinant Human TNFRSF5/CD40 Protein

Catalog No.: RP01218 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 958 P25942-1

Tags

C-hFc&His

Synonyms

CD40;Bp50;CDW40;TNFRSF5;p50; Bp50; CDW40; TNFRSF5

Product Information

Source Purification
HEK293 cells > 97% by SDSPAGE.

Calculated MW Observed MW

45.95 kDa 55-60 kDa

Endotoxin

< 0.1 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μ m filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

6	400-999-6126
\bowtie	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

Background

Basic Information

Description

Recombinant Human TNFRSF5/CD40 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Glu21-Arg193) of human CD40/TNFRSF5 (Accession $\#NP_001241.1$) fused with a Fc, $6\times$ His tag at the C-terminus.

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized Recombinant Human CD40 Ligand at 2 μ g/mL (100 μ L/well) can bind Recombinant Human CD40 with a linear range of 224-898 ng/mL.


Storage

Store at -20° C. Store the lyophilized protein at -20° C to -80° C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20 $^{\circ}$ C for 3 months, at 2-8 $^{\circ}$ C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Validation Data



Recombinant Human TNFRSF5/CD40 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 55-60 kDa.



The purity of human CD40/TNFRSF5 Protein (Cat.RP01218) was greater than 90% as determined by SEC-HPLC.



Immobilized Recombinant Human CD40 Ligand at $2\mu g/mL$ ($100~\mu L/well$) can bind Recombinant Human CD40 with a linear range of 224-898 ng/mL.