

Recombinant Human Nectin-1/PVRL1/CD111 Protein

Catalog No.: RP01219 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	5818	Q15223-1

Tags

C-His

Synonyms

NECTIN1;CD111;CLPED1;ED4;HlgR;HV1S;
HVEC;OFC7;PRR;PRR1;PVRL1;PVRR;PVRR
1;SK-12;nectin-1

Product Information

Source	Purification
HEK293 cells	> 97% by SDS- PAGE.

Endotoxin

< 0.1 EU/μg of the protein by LAL
method.

Formulation

Lyophilized from a 0.22 μm filtered
solution of PBS, pH 7.4. Contact us for
customized product form or formulation.

Reconstitution

Centrifuge the vial before opening.
Reconstitute to a concentration of
0.1-0.5 mg/mL in sterile distilled water.
Avoid vortex or vigorously pipetting the
protein. For long term storage, it is
recommended to add a carrier protein or
stabilizer (e.g. 0.1% BSA, 5% HSA, 10%
FBS or 5% Trehalose), and aliquot the
reconstituted protein solution to
minimize free-thaw cycles.

Background

Basic Information

Description

Recombinant Human Nectin-1/PVRL1/CD111 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Met1-Thr334) of human Nectin-1/PVRL1/CD111 (Accession #NP_002846.3) fused with a 6×His tag at the C-terminus.

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized Recombinant Human Nectin-1 3 μg/mL (100 μL/well) can bind Recombinant Human Nectin-3 with a linear range of 180-720 ng/mL.

Storage

Store the lyophilized protein at -20°C to -80 °C for long term.

After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Contact

 | 400-999-6126


 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn


Validation Data



Active Recombinant Human Nectin-1/PVRL1/CD111 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 50-55 kDa.



The purity of human Nectin-1/PVRL1/CD111 Protein (Cat.RP01219) was greater than 90% as determined by SEC-HPLC.



Immobilized Recombinant Human Nectin-1 3µg/mL (100 µL/well) can bind Recombinant Human Nectin-3 with a linear range of 180-720 ng/mL.