

Catalog No.: RP01230 **Recombinant**

Species	Gene ID	Swiss Prot
Mouse	12519	O00609

C-His

B7;B7-1;B7.1;BB1;CD28LG;CD28LG1;LAB7;CD80;B7;B7-1;B7.1;BB1;CD28LG;CD28LG1;LAB7;CD80

Source	Purification
HEK293 cells	≥ 90 % as determined by SDS-PAGE.

Calculated MW	Observed MW
24.52 kDa	40-55 kDa

< 0.1 EU/μg of the protein by LAL method.

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

 | 400-999-6126

✉ | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

The protein is a membrane receptor that is activated by the binding of CD28 or CTLA-4. The activated protein induces T-cell proliferation and cytokine production. This protein can act as a receptor for adenovirus subgroup B and may play a role in lupus neuropathy.

Recombinant Mouse B7-1/CD80 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Val38-Lys245) of mouse B7-1/CD80 (Accession #NP_033985.3) fused with a 6×His tag at the C-terminus.


Measured by its binding ability in a functional ELISA. Immobilized Recombinant Mouse CD80 at 2 µg/mL (100 µL/well) can bind Recombinant Human CTLA-4 with a linear range of 16-66 ng/mL.

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Mouse B7-1/CD80 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized Recombinant Mouse CD80 at 2 μ g/mL (100 μ L/well) can bind Recombinant Human CTLA-4 with a linear range of 16-66 ng/mL.