

Recombinant Human Follistatin/Follistatin 315/FST Protein

Catalog No.: RP01249 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 10468 P19883

Tags

C-His

Synonyms

FST;FS

Product Information

Source Purification

HEK293 cells \geq 95 % as

determined by SDS-

PAGE.

Calculated MW Observed MW

35.59 kDa 40-50 kDa

Endotoxin

 $< 0.1 \; \text{EU/}\mu\text{g}$ of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

a	400-999-6126
×	cn.market@abclonal.com.cn
	www.abclonal.com.cn

Background

Basic Information

Description

Recombinant Human Follistatin/Follistatin 315/FST Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gly30-Trp344) of human Follistatin (Accession $\#NP_037541.1$) fused with a $6\times$ His tag at the C-terminus.

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized recombinant Human Inhibin beta A at 0.5 μ g/mL (100 μ L/well) can bind recombinant Human FST, the EC₅₀ of Human FST is 7.1 ng/mL.

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20 $^{\circ}$ C for 3 months, at 2-8 $^{\circ}$ C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.

Validation Data



Recombinant Human Follistatin/Follistatin 315/FST Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized recombinant Human Inhibin beta A at 0.5 μ g/mL (100 μ L/well) can bind recombinant Human FST, the EC $_{50}$ of Human FST is 7.1 η g/mL.