Recombinant Human ACE-2 Protein

ABclonal www.abclonal.com

Catalog No.: RP01277 Recombinant 1 Publications

Sequence Information

Species Gene ID **Swiss Prot** Human 59272 O9BYF1

Tags

N-His

Synonyms

ACE2; ACEH; angiotensin-converting enzyme 2;ACEH

Product Information

Purification

HEK293 cells

≥ 95 % as determined by SDS-

PAGE.

Calculated MW Observed MW

84.44 kDa 110-120 kDa

Endotoxin

< 0.1 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

<u>a</u>		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

Background

Basic Information

Description

Recombinant Human ACE-2 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gln18-Ser740) of human ACE2 (Accession #Q9BYF1) fused with a 6×His tag at the N-terminus.

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized recombinant Human ACE2 at 2 µg/mL (100 µL/well) can bind recombinant SARS-COV-2 S1 Protein with a linear range of 0.49-9.31 ng/mL.


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human ACE-2 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized recombinant Human ACE2-His at 2 μ g/mL (100 μ L/well) can bind recombinant SARS-COV-2 S1-mFc Protein with a linear range of 0.49-9.31 ng/mL.