

Recombinant Human Apolipoprotein A-II/APOA2 Protein

Catalog No.: RP01441 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 336 P02652

Tags C-hFc&Avi

Synonyms

APOA2;Apo-AII;ApoA-II;apoAII; ApoA-II; apoAII

Product Information

Source Purification HEK293 cells > 95% by SDS-PAGE.

Endotoxin

 $< 0.1 EU/\mu g$

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

6		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

Background

Apolipoprotein A-II(Apo-AII for short), also known as Apolipoprotein A2, is a secreted protein which belongs to the apolipoprotein A2 family. It exists as a disulfide-linked homodimer; and also can form a disulfide- linked heterodimer with APOD. APOA2 is the 2nd most abundant protein of the high density lipoprotein particles. This protein may stabilize HDL (high density lipoprotein) structure by its association with lipids, and affect the HDL metabolism. Defects in APOA2 gene might cause apolipoprotein A-II deficiency or hypercholesterolemia.

Basic Information

Description

Recombinant Human Apolipoprotein A-II/APOA2 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ala19-Gln100) of human Apolipoprotein A-II/APOA2 (Accession #NP_001634.1) fused with a Fc, Avi tag at the C-terminus.


Bio-Activity

Storage

Store the lyophilized protein at -20°C to -80°C for long term. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human Apolipoprotein A-II/APOA2 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 45-50kDa.