

Recombinant Human Mature TGF-beta 1 Protein

Catalog No.: RP01458 Recombinant 9 Publications

Sequence Information

Species **Gene ID** **Swiss Prot**
 Human 7040 P01137

Tags

No tag

Synonyms

TGFB1; CED; DPD1; LAP; TGFB; TGFbeta; transforming growth factor beta-1;TGF-beta 1;CED;DPD1;LAP;TGFB;TGFbeta;TGF-β

Product Information

Source **Purification**
 HEK293 cells $\geq 95\%$ as determined by SDS-PAGE.

Calculated MW **Observed MW**
 12.79 kDa 15 kDa

Endotoxin

< 0.01 EU/ μ g of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μ m filtered solution of 50 mM Glycine, 150 mM NaCl, pH 3.5.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

Contact

	400-999-6126
	cn.market@abclonal.com.cn
	www.abclonal.com.cn

Background

TGF-beta 1 is a member of the transforming growth factor beta (TGF-beta) family. The transforming growth factor-beta family of polypeptides are involved in the regulation of cellular processes, including cell division, differentiation, motility, adhesion and death. TGF-beta 1 positively and negatively regulates many other growth factors. It inhibits the secretion and activity of many other cytokines including interferon- γ , tumor necrosis factor-alpha and various interleukins. It can also decrease the expression levels of cytokine receptors. Meanwhile, TGF-beta 1 also increases the expression of certain cytokines in T cells and promotes their proliferation, particularly if the cells are immature. TGF-beta 1 also inhibits proliferation and stimulates apoptosis of B cells, and plays a role in controlling the expression of antibody, transferrin and MHC class II proteins on immature and mature B cells. As for myeloid cells, TGF-beta 1 can inhibit their proliferation and prevent their production of reactive oxygen and nitrogen intermediates. However, as with other cell types, TGF-beta 1 also has the opposite effect on cells of myeloid origin. TGF-beta 1 is a multifunctional protein that controls proliferation, differentiation and other functions in many cell types. It plays an important role in bone remodeling as it is a potent stimulator of osteoblastic bone formation, causing chemotaxis, proliferation and differentiation in committed osteoblasts. Once cells lose their sensitivity to TGF-beta 1-mediated growth inhibition, autocrine TGF-beta signaling can promote tumorigenesis. Elevated levels of TGF-beta 1 are often observed in advanced carcinomas, and have been correlated with increased tumor invasiveness and disease progression.

Basic Information

Description

Recombinant Human Mature TGF-beta 1 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ala279-Ser390) of human Mature TGF-beta 1 (Accession #NP_000651.3) fused with no tag.

Bio-Activity

1. Measured by its binding ability in a functional ELISA. Immobilized Human TGF-beta 1 at 2 μ g/mL (100 μ L/well) can bind Human TGFBR2 with a linear range of 0.78-11 ng/mL. 2. Measured by its ability to inhibit the IL-4 (Catalog: RP01161)-dependent proliferation of HT-2 mouse T cells. The ED₅₀ for this effect is 12.85-51.40 pg/mL, corresponding to a specific activity of 1.95×10^7 ~ 7.78×10^7 units/mg.

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.


Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.


Validation Data



Recombinant Human Mature TGF-beta 1 Protein was determined by SDS-PAGE under reducing (R) and non-reducing (NR) conditions.



Immobilized Human TGF-beta 1 at 2 μ g/mL (100 μ L/well) can bind Human TGFBR2 with a linear range of 0.78-11 ng/mL.



Recombinant Human Mature TGF-beta 1 inhibit the IL-4(Catalog: RP01161)-dependent proliferation of HT-2 mouse T cells. The ED₅₀ for this effect is 12.85-51.40 pg/mL, corresponding to a specific activity of $1.95 \times 10^7 \sim 7.78 \times 10^7$ units/mg.